

#### Acknowledgements:

Mr. Murat Akyol	Miss Jen Lumsdaine
Ms Jackie Bradie	Mrs. Shelagh Salter
Dr. Conal Daly	Mrs. Betty Sloan
Sr. Caroline Stephenson	Miss Lynne Thomson
Prof John Forsythe	Mrs. Gillian Walker
Miss Clare Hodgkinson	Ms Lorna Marson
Dr. Ellon McGregor	Mrs Laura Pairman

Designed by Stuart Brett, The Graphics Lab,  
The University of Edinburgh  
© 2013 NHS Lothian

# Kidney Transplant

Your questions answered



# C O N T E N T S

Page

02

**Section 1** - Introduction and assessment for transplantation

10

**Section 2** - The waiting period

12

**Section 3** - The operation

17

**Section 4** - Results and complications

22

**Section 5** - Medications

25

**Section 6** - Living with a transplant





## INTRODUCTION

### Welcome to the Renal Transplant Unit at the Royal Infirmary of Edinburgh

The first kidney transplant performed in the Royal Infirmary was in 1961 and indeed this was the first successful kidney transplant in the United Kingdom. Since that time over 1,000 patients have experienced the benefits of kidney transplantation in the Edinburgh Renal Transplant Unit.

On 5th June 1995 the Renal Transplant Unit joined with the Scottish Liver Transplant Unit, so that the efforts of the transplant surgeons, doctors, nurses and other staff could be combined.

Within this booklet we provide information for patients awaiting kidney transplantation, those about to receive a transplant and families and friends of renal patients. We are very keen that patients get a chance to visit the Unit long before a transplant is performed, so that they may ask any questions or tell us of worries that they may have.

The information in this booklet is a basic guide only. Please contact us if you wish to know any more.

#### Useful contact numbers:

0131 536 1000	RIE Switchboard
0131 242 2068/9	Ward 206 Transplant Unit. Patient enquiries/out of hrs contact number
0131 242 1171/7	Ward 117 Transplant High Dependency Unit
0131 242 1151	Ward 115 Renal High Dependency Unit
0131 242 7850	Renal Social Workers
0131 242 1635	Transplant Co-ordinator Secretary

#### Where are the Outpatient Clinics?

The transplant assessment and follow-up clinics are held in Outpatients Department 1, Royal Infirmary of Edinburgh (OPD1 RIE). This clinic is located on the ground floor and can be accessed via the main entrance to the hospital.



### The Transplant Unit Staff (Royal Infirmary of Edinburgh)

Consultant Surgeons		Prof John LR Forsythe Prof Steve Wigmore Mr Murat Akyol Mr John Casey Miss Lorna Marson Mr Gabriel Oniscu Mr James Powell Mr Ian Currie Ms Anya Adair
Consultant Physicians		Dr Caroline Whitworth Dr Simon Watson Dr Paddy Gibson Prof Neil Turner Dr Richard Phelps Dr Jeremy Hughes Dr Jane Goddard Dr David Kluth Dr Wendy Metcalfe Dr John Neary Dr Lorna Henderson
Live Donor Transplant Co-ordinators	0131 242 1703	Ms Jen Lumsdaine Mrs Sarah Lundie Mrs Yvonne Perry
Renal Transplant Co-ordinators	0131 242 1728 0131 242 1727 0131 242 1727	Mrs Laura Pairman Miss Alison Glover Miss Debbie Mills
Pancreas and Islet Co-ordinators	0131 242 1704 0131 242 1730	Mrs Christine Jansen Mrs Kirsty Duncan Mrs Melanie Phillips
Senior Nursing Staff		C/N Jacqueline Reid C/N Steve Elliott

### What happens when a kidney becomes available?

When a kidney becomes available, someone from the transplant team will contact you and ask you to make your way to the Transplant Unit at the RIE. This call may happen at any hour of the day or night.

Remember that the hospital entrances are locked from 9pm to 6am. There are buzzers at the entrances where the security staff will give you access to the hospital. If you are carrying dialysis equipment, we suggest you enter via the Accident and Emergency Department, as a porter can be called to help you.

#### You will be asked:

1. If you are fit and well
2. Where and when you were last dialysed
3. To have nothing more to eat or drink.



## Is there car parking at the hospital or can I get there by bus?

### *By Car:*

The Royal Infirmary of Edinburgh is on Old Dalkeith Road (A7) on the south side of Edinburgh.

### *Directions from City Centre:*

Leave the city centre on North Bridge at the east end of Princes Street. Follow this road, taking the first exit (left) at the roundabout with Liberton Road and Lady Road. Go straight over the Lady Road roundabout and take the third exit (right) off Cameron Toll roundabout on to Old Dalkeith Road (A7). Turn left into the hospital after three-quarters of a mile.

### *Directions from Bypass:*

Leave the Edinburgh City Bypass (A720) at the Sheriffhall roundabout on to Old Dalkeith Road (A7) toward Edinburgh City Centre. Turn right into the hospital after just over 2 miles.

Entry to the site is via two access roads, one to the north and one to the south of the site. The north access road leads to the patient and visitor car park. There are free disabled parking spaces located near the main entrances. The current parking charges for all other patients and visitors are:

0-1 hours	£ 1.30
1-2 hours	£ 2.30
3-4 hours	£ 3.30
4-5 hours	£ 5.30
5-6 hours	£ 6.50
Over 7 hours	£7.00

The nearest entrance and drop off point to our department is by entering the North Junction from Old Dalkeith Road and going to Entrance 1.

### *By Public Transport:*

The following services stop at the hospital either at the west main entrance bus stops, or close by on Old Dalkeith Road.

LRT 7, Newhaven to RIE  
 LRT 8, Muirhouse to RIE  
 LRT 18, Gyle to RIE  
 LRT 21, Gyle to RIE  
 LRT 24, West Granton to RIE  
 LRT 33, Baberton Mains to Sheriffhall Park and Ride  
 LRT 38, Granton to RIE  
 First 86, Charlotte Square to Mayfield via RIE  
 First 86A, Charlotte Square to Birkenside via RIE  
 First 18, Gyle Centre to RIE  
 First 328, Coysland/Dalkeith Campus to Musselburgh  
 First 428, Coysland/Dalkeith Campus to Musselburgh

For full timetables or further information, contact:

Lothian Buses: 0131-554 4494, [www.lothianbuses.co.uk](http://www.lothianbuses.co.uk)

First Edinburgh: 0871 2002233, [www.firstedinburgh.co.uk](http://www.firstedinburgh.co.uk)

Traveline: 0871 2002233

### *By Foot / Cycle:*

New cycle/bus lane from Cameron Toll to the Royal Infirmary of Edinburgh via Old Dalkeith Road, as well as improvements to the cycle and footpaths on Craigmillar Castle Road, may make this an option for some patients and visitors. There is currently provision to park both bicycles and motorbikes on site, free of charge. Motorcycle parking is within marked spaces outwith the car parks at various points around the perimeter of the building. Cycle parking is available close to all the entrances to the building, in the form of Sheffield stands. CCTV monitors these areas at all times.

### Do I have to pay to park at the hospital?

Unfortunately the parking at the hospital is run by an external business and we have no control over the parking fees however, we understand that for some people this may be very expensive and also that you have travelled a distance to the hospital. When a patient has been called in for transplant, the transplant staff have guidelines on issuing passes, which will allow parking at a reduced fee. Please discuss this with the nurse in charge of the unit at this time.

For clinic appointments, you may be able to claim this back from your health authority.

### Who will I meet?

Some of those responsible for your care include:

**Doctors** - a team of surgeons, physicians, anaesthetists and radiologists will look after you. You may already know some of them from outpatient clinics or previous hospital admissions. As the Royal Infirmary of Edinburgh is a Teaching Hospital, you will come across junior as well as senior doctors and, perhaps, medical students.

**Nurses** - During your operation you will be looked after by theatre staff. On your return to the Transplant Unit your care will be planned and carried out by nursing staff who work in teams, as this ensures the best possible standard of care for our patients.

**Transplant Co-ordinators** - You should have met at least one of the transplant co-ordinators before your admission. As they are based in the Transplant Unit, you will be sure to meet them again during your hospital stay.

**Chaplain** - The Royal Infirmary has a number of chaplains of different denominations who are on hand to offer support to any patient or their relatives in a general way, while offering religious and spiritual help. The chaplains can also arrange support for those of our patients who are not members of the Christian faith. Please do not hesitate to ask for a visit if they can be of any help to you or your family.

### What are the benefits of kidney transplantation?

Kidney transplantation is an excellent treatment for kidney failure and offers the chance to lead a nearly normal life with:

- Freedom from dialysis
- Improved capacity to socialise and enjoy life to the full
- Return of strength and energy
- Freedom from dietary restrictions
- No restriction on physical activity (except contact sports)

### What are the disadvantages of kidney transplantation?

- Risk of failure
- Infection
- Malignancy
- Diabetes
- Side-effects of drugs necessary for transplant

*See section on complications for further details.*

### Assessment for transplantation

The first step in being considered for transplant is to talk to your Renal Doctor who will refer you for assessment if appropriate.

### Does everyone with kidney failure get a transplant?

No. Only about 50% of patients with renal failure are on the active transplant list. Each patient will be assessed individually to see if transplantation is the best option.

### Reasons why kidney transplantation may not be the best treatment option:

- History of recent cancer
- Major surgery
- Bad heart, lung or blood vessel disease.

If you do not wish to be on the transplant list, that is your right.

### Why do I have to see the transplant surgeon?

This appointment is to assess your health and fitness to ensure you do not require any further investigations before your transplant operation.

### What will happen at the visit?

The transplant surgeon will take this opportunity to inform you about the transplant operation and introduce you to the transplant co-ordinator. There will be a discussion with you about your general health. The transplant surgeon will decide if you require to undertake any tests before going onto the transplant list. The visit to the clinic also allows you to ask any questions about kidney transplants, so that you can decide whether transplant is the right thing for you.

You may find it useful to bring along a relative or friend for support and to discuss any issues raised at your consultation. If you already have any questions that you would like to ask, it may be an idea to write them down and bring them with you on the day.

### Are there any risks from having a transplant?

All organ donors are assessed to find out if they have any infection or illness that could be passed to potential transplant recipients. Every possible precaution is taken to prevent the transmission of viral infections, cancer or any other potentially life threatening condition to patients who receive an organ for transplantation. Whilst every effort is made to gather this information, we are unable to give a 100% guarantee that the organ that you will receive is completely free of risk factors.

We are very careful in assessing the function of the donor organ before transplant. Due to the shortage of organ donors, we now consider some people to be potential donors who we would not have accepted as donors a few years ago. This means you could receive an organ from an older donor or that the organ to be transplanted is not functioning perfectly. The results of transplants from donors like these are still seen to be very acceptable. Therefore it is felt that the balance of risk is in favour of going ahead with such a transplant rather than leaving a patient waiting on the list.

If there are any special risk factors associated with a donor over and above those described, they would be discussed with you immediately before any transplant procedure.

## THE WAITING PERIOD

### What should I do while waiting for my transplant?

- Stop smoking
- Continue to follow dietary advice
- Keep to your fluid restriction
- Take the medicines prescribed by your doctors
- Exercise
- Try not to restrict your lifestyle
- Continue to attend your dentist regularly
- If overweight, consider dietary advice/weight loss.

All the above will help to keep you in good health for when you are called for your transplant operation. Also:

- Tell your transplant co-ordinator if you are going on holiday, as they need to suspend you from the list while you are away
- Inform your transplant co-ordinator of any change of address/ telephone number.

### Can you tell me about the waiting time?

After you have made the decision to have a kidney transplant, there will be a period of waiting until a suitable donor kidney becomes available - unfortunately we cannot predict how long this will be.

If at any time you are unwell or in hospital, it is likely that you will be temporarily suspended from this list. If you require further clarification of this, please contact your renal physician.

Organ Donation and Transplantation (ODT) allocates kidneys by blood group and tissue type. This system is in place to ensure the best match between you and the donor kidney and the average waiting time is two and a half to three years. This can be a difficult time for both you and your family. You will be reviewed on a regular basis whilst on the Transplant List.

### Where do donor organs come from?

There are two sources of kidneys for transplant:

- 1 From a living donor
- 2 From a person who has died and the family have agreed to organ donation: either people who have suffered brain death, usually as a result of an injury or brain haemorrhage, known as Donation after Brain Death (DBD) or Donation after Circulatory Death (DCD) from patients who are declared dead following absence of breathing or heartbeat. The organs are retrieved rapidly following this type of donation.

### What are the potential advantages of the DCD programme?

It will increase the likelihood of you receiving a kidney transplant; DCD kidneys will only be used for patients on our waiting list.

### What are the potential disadvantages of receiving a DCD kidney?

After transplant, kidneys may have delayed graft function (a period where the kidney "sleeps" until recovery). This is more common if the kidney comes from DCD but does not affect the long-term function of the transplant. At the time we call you in for transplant we will inform you if your kidney is DCD.

### Can I have any information about my donor?

It is possible to find out some details about the donor of your kidney and write a letter to the donor family. Your transplant co-ordinator will discuss this with you at an appropriate time after the transplant has taken place.

### Will I be asked to take part in any clinical research trials?

The field of transplantation is constantly evolving. You may be asked to take part in clinical research trials. Any trial will be discussed with you in full. If you decide not to take part in any trial your treatment will not be affected in any way.



## THE OPERATION

### What do I need to consider when called for my transplant operation?

- How are you going to get to hospital?
- Who is going to look after your children/pets?
- Things you need in hospital: nightclothes, wash things, and a little money for papers/ telephone
- Lock up the house remembering to turn off all appliances
- Inform someone where you are, pop a note through your neighbour's letterbox; you may need them to cancel your milk and papers
- When we call you, tell us if you are unwell: we require you to be in the best possible health for your transplant operation
- You may be asked to bring your dialysis equipment, e.g. Homechoice machine, as it may be required before the transplant. You do not need to bring fluids
- Bring all your usual medications with you to the hospital.



### Is it definite that I will have the transplant operation at this time?

No - not until the doctors have assessed you to ensure you are fit enough for surgery. We may also need to do a final crossmatch test between you and the donor. In addition, the surgeon will need to examine the kidney carefully to ensure there is no problem to prevent the transplant going ahead.

### What do I do when I arrive at the transplant unit/renal ward?

Report to the nurses in the duty room of the transplant unit/ renal ward. You will be shown to your bed and the nurses will inform the medical staff of your arrival. You will require some final tests to ensure you are fit and to see if you require dialysis before surgery.

### Where do I have my operation?

The transplant operation takes place in the operating theatre where you will have a general anaesthetic. The operation takes between 2-3 hours.

### Where does the kidney go?

The transplanted kidney is placed in the right or left groin area. It is placed here as it is close to your bladder. The operation requires the surgeon to join up your blood vessels to those of the kidney. Also the urine tube (ureter) from the kidney needs to be joined to your bladder.

### What happens to my own kidneys?

Your own kidneys are not normally removed unless there is a reason to do so. You will end up with three kidneys.



### What tubes will I have after the operation?

- You will have a tube placed into your bladder (a urinary catheter). This is to measure the amount of urine your transplanted kidney is producing. It will be removed after five days, when the doctors are happy that your kidney is functioning well and you are drinking more fluid than you are passing.
- You may have a tube (drain) from the operation site, where your kidney has been placed. This will be removed within a few days. This is to take away any blood or fluid from the operation site to prevent any swelling or infection. Not all patients need to have a drain.
- You will have a plastic tube (a stent) placed inside the urine tube (ureter) that connects the kidney to your bladder. This prevents any kinking of the ureter after the operation. The stent needs to be removed about six to eight weeks after your transplant operation, so you will need to come back for a day stay in hospital and a trip to the operating theatre. Your peritoneal catheter may be removed at the same time.
- You will have a drip line in your hand or arm and also a central line (neckline).
- You will be given control of your analgesia (painkiller). This is attached to one of your drip lines and a button control will be placed in your hand. When you press the button you will deliver a painkiller that works very quickly to prevent any further discomfort. This is called PCA - patient controlled analgesia.

### Will the kidney work straight away?

After the transplant operation, the kidney may work well from the start, but it is quite possible that it will take a little time for it to work efficiently. Often the kidney reacts to the transplant process by a period of poor function - it 'goes to sleep'. During this time dialysis may need to be continued. This period of poor function should not cause alarm because in nearly every case, the kidney gradually recovers to work well, allowing dialysis to stop. If you are on peritoneal dialysis you may need a short period of haemodialysis.

## The Recovery Period

### Will I need dialysis after the transplant?

In most cases you will not require dialysis. A small number of patients may require to continue or start dialysis until such time as their kidney begins to function.

### Will I get rejection?

Rejection is a normal reaction of your body to tissue that it detects as foreign. Rejection is common and affects up to 25% of patients after transplant. In almost all cases of rejection, drugs are effective at treating it.

### Will I need a transplant biopsy?

It is very common to have a transplant biopsy. This may help to explain why the kidney is not functioning.



The biopsy is done using ultrasound. Local anaesthetic is given at the site of the biopsy and a special needle device is used to obtain a small piece of kidney tissue. You may have had a biopsy of your own kidney. Transplant biopsy is very much more straightforward than this.

### How soon can I eat after my operation?

You will not be able to eat or drink immediately after the transplant operation. After a few hours you should be able to start drinking sips of water and this will gradually be built up to free fluids and a light diet. Within a couple of days you should be eating normally.

Your body's requirements for energy (calories) and proteins are high after the stress of an operation. The dietitian will assess your diet on the ward and will provide advice if your appetite is poor.

The doctor will assess how much fluid you should drink every day. Remember, it is often necessary to drink much more fluid than you have been used to on dialysis - it may be many litres each day.

Some people may pass large amounts of urine (polyuria) after the transplant operation. This is nothing to worry about and will eventually settle down. However, you may need to have some of your fluids through a drip at this time.

### How soon will I be up and around after my operation?

In order that you make as full a recovery as possible after your operation, you will receive a visit from the unit's physiotherapist. The physiotherapist's role is to teach you a range of simple exercises that will help you avoid the complications that can occur following surgery, e.g. the risk of chest infection can be reduced by regular deep breathing and coughing exercises.

It is also important that you start gentle mobilisation in the first few days post-op. This has many benefits and again will help to minimise the risks associated with surgery. The physiotherapist and nursing staff will assist you with this.

The physiotherapist will continue to see you for a few days after your operation to ensure your lungs are clear and that you are up and about without any issues. Before you are discharged home it is also useful to discuss safe progression of exercise and the re-introduction of any hobbies in the longer term.

## RESULTS AND COMPLICATIONS

### How successful is kidney transplantation?

1-year DBD transplant survival - 94% (2006 - 2009)

1-year DCD transplant survival - 92% (2006 - 2009)

5-year DBD transplant survival - 84% (2003 -2005)

5-year DCD transplant survival - 86% (2003 - 2005)

10-year DBD transplant survival - 69% (1997 - 1999)

10-year DCD transplant survival - 70% (1997 - 1999)

1st adult cadaveric renal transplant UK.

Figures provided by Organ Donation and Transplantation.

In general terms, kidney transplantation is a very successful treatment for renal failure. However complications can occur and some of these are listed below.

### Early complications can include:

**Delayed Function** - not all transplanted kidneys will work straight after the operation. This is not uncommon and occurs in at least 20% of transplants. While waiting for the new kidney to start working, you will probably require dialysis. Sometimes a patient treated by peritoneal dialysis can continue this treatment after transplant. More commonly haemodialysis is necessary. If you do not have a fistula, a special dialysis catheter can be put into a vein in your neck at the time of your operation.

**Rejection** - your body will recognise that your new kidney is 'foreign' and the natural response is to attack and reject it. You will be given drugs to reduce the chance of this happening. Despite these drugs, early rejection is fairly common, especially in the first three months following your transplant. The signs of rejection can be:

- Pain over your new kidney
- Fever
- Reduced urine output
- Flu-like symptoms.

However, in many cases there are no signs of rejection and it is detected following one of your regular blood tests. Most episodes of rejection at this stage will respond to treatment. This will require an increased dose of anti-rejection drugs and/or a change to your medication. Very rarely the rejection cannot be controlled. In these cases the kidney will need to be removed and you will have to return to dialysis. When you feel well enough you will be given the opportunity to discuss going back on to the transplant waiting list.

**Infection** - the anti-rejection treatment that you must take following your transplant means that you will be more likely to develop infections. The risk is greater in the early stages after your transplant when the doses of the drugs are higher. It is important that you contact the transplant unit quickly if you think you have an infection. If you know of any friends or relatives who have a serious infection, it is a good idea if you avoid visiting them while they are unwell.

**Thrombosis** - this is a rare complication following transplant, when a clot forms in the vein or the artery of the new kidney. This means that the blood supply to the kidney will stop. If this happens you will have to go back to theatre and in most cases the kidney will have to be removed.

**Urine Leak** - occasionally a leak will occur where the ureter (urine drainage tube) from the transplanted kidney joins your bladder. This usually requires a return to theatre and an operation to have the leak repaired.

**Bleeding** - as with all major surgery there is a risk of internal bleeding. This may require a blood transfusion and a further operation to stop the bleeding.

**Deep Venous Thrombosis (DVT)** - following all surgery there is a risk of clots forming in the veins of the leg. This problem is less common in patients with renal disease but can still occur. Under certain circumstances these clots can be dangerous. To prevent this happening you will be given subcutaneous (under the skin) injections of heparin for the first few days following the transplant operation.

## Late complications:

**Weight gain** is a common problem after successful renal transplantation because of improved appetite with better renal function and steroid treatment. It can be minimised by eating a healthy diet (see section on diet) and by taking regular exercise.

**High Blood Pressure** is extremely common following kidney transplantation because of the immunosuppressive drugs and pre-existing kidney disease. Many people require blood pressure-lowering drugs long-term. Rarely, high blood pressure can be caused by narrowing in the transplant artery.

**Diabetes** can develop for the first time following renal transplantation as a consequence of drug therapy (particularly tacrolimus and high dose steroid treatment) in up to one in six patients. It is generally treated by means of a low sugar diet, but tablet treatment and sometimes insulin injections are necessary.

**Cholesterol levels** and other blood fats can increase after a kidney transplant, principally as a side-effect of drugs, with various other causes contributing. A healthy diet will help, though cholesterol-lowering treatment might also be required.

**Cardiovascular Diseases** such as angina, heart attacks, circulatory problems and strokes are all more common in transplant and dialysis patients than in people without renal disease. Receiving a kidney transplant does not reduce the risk, and it is therefore most important to stop smoking and to follow any advice on blood pressure, cholesterol and diabetes. Keeping your weight down and taking exercise regularly are also sensible measures.

**Osteoporosis** and other disorders of bone are more frequently found in both men and women after renal transplantation. This is sometimes related to high doses of steroid, though a number of different factors can contribute. If you have already had a fracture, or are female and post-menopausal, you may be advised to have additional drug treatment such as hormone replacement. Preventative medication might also be recommended around the time of your operation.

**Liver disease**, mainly inflammation, can sometimes follow kidney transplant surgery as a consequence of viral infections or drug treatment. In this country it rarely causes major problems.

**Recurrence of the original disease that damaged your own kidneys** is a recognised complication but an unusual cause of transplant failure. Some types of renal disease are more likely to recur than others and you will be monitored closely for this problem. Occasionally a new form of nephritis can develop in a transplant kidney for reasons that are not clear.

**Cancer** is more common in the longer term in people with kidney transplants because of immunosuppressive treatment and types of viral infection. One of the commonest types of tumour to develop is skin cancer. Careful avoidance of sun exposure and use of powerful sunscreens help reduce the risk to skin. Other types of tumours such as lymphomas are seen more frequently than normal, while some types are less common.

It is clear that many of these side-effects can be attributed to the immunosuppressive medication you are required to take. Unfortunately without these drugs, the kidney transplant will fail. However, new drugs with fewer unwanted effects are continually under development and your therapy might change in the future if these become widely available and accepted.

**Chronic Rejection** is a slow process and occurs over months or years, with a slow worsening of kidney function tests. It is also relatively common but no specific treatment has yet been shown to stop this form of rejection.

## Is infection more common after a transplant?

The operation itself and your immunosuppressant drugs make you more susceptible to infections. The majority of these infections, such as wound, urinary and chest infections are caused by relatively common bacteria, which are usually easily treated with relatively common antibiotics. However, because of your immunosuppressant drugs you are also susceptible to serious and occasionally life-threatening diseases from rarer infections such as Cytomegalovirus (CMV), fungal infections and pneumocystis jirovecii.



CMV Disease deserves special mention. In patients who are not immunosuppressed by drugs or an illness, CMV is usually a very mild viral illness. However, in patients on immunosuppressant drugs it may be much more serious and may be life-threatening. Patients may feel vaguely unwell, feverish or breathless. Patients at greatest risk of developing serious CMV Disease are those who have never had CMV before but receive a kidney from a donor who has had CMV. However, with present drug regimens, serious infection with these organisms can usually be prevented (you may be on additional drugs for this) or if they do occur can nearly always be treated. With time, the increased risk of infection because of immunosuppressants lessens but never goes away completely.



## MEDICATIONS

### How are my tablets going to change?

Some medications will be stopped. Some of the medications you are currently taking are needed because your kidney is not working properly. After a transplant, these medicines may be stopped. They include phosphate binders, anti-itch tablets, iron supplements and erythropoietin.

**Some medications will be started.** The most important new tablets you will be started on are called IMMUNOSUPPRESSANTS. The immune system is our body's natural defence mechanism. It is programmed to recognise and destroy anything unfamiliar. This includes a transplanted kidney as well as the organisms that cause infection.

After a kidney transplant, you will be prescribed immunosuppressants to prevent the immune system from rejecting the new kidney. You must take them every day while your kidney transplant is functioning.

### What immunosuppressants will I take?

You are likely to be on the following medicines:

1. Tacrolimus (Prograf or Advagraf)
2. Prednisolone
3. Mycophenolate Mofetil (Cellcept or Myphenax)
4. Mycophenolate Sodium (Myfortic)

**If you do not take these medications as prescribed, it will result in rejection and the loss of your new kidney.**

These drugs need to be finely tuned, as too little may lead to rejection and too much may lead to infection. Early after the transplant the need for immunosuppression is high but this need lessens with time, so these drugs will be slowly reduced by your doctors.

One further drug, which is licensed for renal transplantation, is Basiliximab (Simulect). This drug (known as a monoclonal antibody) is given in the early period after a transplant via a drip or infusion.

## What are the possible side-effects?

Immunosuppressants, like most medicines, can have some unwanted side-effects. However, just because a medicine has the potential to cause adverse effects, it does not necessarily mean you will get them. All anti-rejection drugs will increase your risk of infection. The risk lessens as they are cut down, but common-sense precautions will be necessary. Any signs of serious infection will need to be brought to the attention of the transplant unit/ renal ward immediately.

The individual immunosuppressives may have their own side-effects as follows:

**Tacrolimus** - shaking of the hands, headaches, impaired vision, 'pins and needles', raised blood sugar levels (diabetes). Kidney damage at high blood levels.

**Prednisolone** - irritation of the gut lining, indigestion, increase in appetite, weight gain, rounded face, thinning skin and bones and raised blood sugar levels (diabetes).

**Mycophenolate Mofetil** - diarrhoea, bloating, heartburn, nausea and vomiting.

**Mycophenolate sodium** – side-effects are similar to those detailed above for mycophenolate mofetil.

Inform the doctors if you experience side-effects - they may want to cut down the dose or give you an alternative.

## What other medications do I have to take?

You may be asked to take the following:

- co-trimoxazole - an antibiotic to reduce the risk of bacterial infection
- ranitidine - to protect the stomach against ulcers and heartburn
- antiviral medications such as valganciclovir
- heparin - to prevent blood clots (for the first few days after the operation).

## What is the self medication programme?

After your transplant, when you are ready, you will have the opportunity to participate in the self-medication programme that operates in the Transplant Unit.

The programme is designed to give you practical information about your medicines and how to take them whilst you are still in hospital. The nursing staff and pharmacist will ensure that by the time you are ready to go home, you feel confident that you know all about your medications and their possible side-effects.

## Can I take any other medicines?

Some of these medicines are affected by and do themselves affect other medicines. Please discuss your current medication with the pharmacist if you need to buy 'over the counter' medicines for minor ailments.

Your GP may wish to contact the transplant unit/ renal ward before giving you any new prescription medicines. If you have any concerns about any aspect of your medicines, please do not hesitate to ask.

If you are taking tacrolimus, the following drugs may interact. Your renal/ transplant doctor should be informed before these drugs are used.

For example:

- erythromycin
- clarithromycin
- fluconazole
- ketoconazole
- St. johns wort
- herbal medicines
- grapefruit juice

Many other drugs also interact. If you or your doctor are unsure, please discuss with your renal/ transplant doctor.

## Herbal medicines

It is suggested that you do not take any herbal medicines (in particular St. johns wort should not be used) without first checking with your renal/ transplant doctor.

## Grapefruit juice

Grapefruit juice taken along with tacrolimus can alter the levels of these drugs in your blood. It is best to avoid grapefruit juice. For further advice speak to your renal/ transplant doctor.

# LIVING WITH A TRANSPLANT

## What about my diet when I go home?

You should not need a special diet if your new kidney is functioning well but this is a good time to think about your general health and the positive things you can do to improve it.

Many common health problems such as heart disease can be linked to a poor diet and lifestyle. Healthy eating is an important part of keeping fit and is one of the positive things you can do to improve your health. Healthy eating is not a 'diet' - it requires a gradual change in your eating behaviour and should become a regular part of your lifestyle.

Healthy eating involves:

- Enjoying what you eat and having a varied diet.
- Eating more fruit and vegetables
- Cutting down on fat, sugar and salt
- Being a healthy weight. Limiting alcohol to below recommended weekly limits (21 units for men, 14 for women)

The transplant co-ordinator will give you more information about healthy eating and food safety before you are discharged home.

## Will I gain weight after my transplant?

Many people gain weight following a transplant, especially in the first year. This can be due to freedom from dietary restrictions, feeling well, increased appetite and lack of exercise. This is not helped by steroids, which are necessary after a transplant.

Maintaining a healthy weight is very important for your health, as being overweight puts a strain on your body, contributing to many health problems such as high blood pressure, heart disease and diabetes.

If you are overweight, it is worthwhile trying to lose some weight before you receive your transplant. If you wish to discuss this, please ask to speak to your dietitian.

It is possible to maintain a healthy weight after your transplant if you make the necessary changes to your diet at an early stage. Remember - it is much easier to prevent weight gain than it is to try to lose weight and it is sensible to make changes to your eating habits as early as possible to prevent weight gain.

Advice on how to achieve and maintain a healthy weight will be tailored to your individual needs and can be discussed with the dietitian before you are discharged.

**Follow up** - You will need to be seen regularly in the outpatient clinic following your discharge home from hospital. The time between appointments increases depending on how well you are. At these appointments you will have your bloods monitored and be seen by one of the renal /transplant follow-up staff. Do not take your morning tacrolimus tablets on the morning of your clinic visit until after your blood test. Then it should be taken as normal.

If we need to make any changes to your medications/ treatment we will contact you by phone or speak with/ write to your GP. In some cases we may ask you to come into hospital.

**Returning to work** - The aim of a transplant is to allow each recipient to return to an independent lifestyle with a good quality of life. Most transplant recipients remain off work for 2-3 months, but this depends on the individual patient and their occupation. Please discuss when you can return to work with the surgeon or physician caring for you.

**Driving** - Usually you are able to drive at about six weeks following your operation. You should also check with your individual insurance company. Remember always to use your seat belt.

**Equipment** - If your transplant is successful, you should have no further use of your dialysis equipment. Contact your local community dialysis team to arrange for this equipment to be uplifted and for future deliveries to be cancelled.

## Family Planning

**Females** - It is not advisable to consider pregnancy within the first year post transplantation. Transplant patients of child-bearing age should practice birth control following transplantation. Should you wish to take the oral contraceptive pill, discuss this at your outpatient appointment.

**Males** - It is important that our male patients take 'pregnancy advice' regarding fathering a child. We do not advise that you father a child whilst taking certain medications.

**Libido** - Some patients have a change in their libido. This is not uncommon and it may take some time for your sex drive to return to normal. If matters don't improve, please discuss this with a member of the medical staff at your outpatient appointment.

**Sex** - You can resume sexual activity once you feel well enough. However you may find that your sex drive is reduced during the early convalescence period, although this will gradually return to normal.

**Impotence** - Some patients may experience/ continue to experience impotence. This is not uncommon and may in some cases be easily treated. Please discuss any problems you may be experiencing with the medical staff at your outpatient appointment.

**Pregnancy** - It is not advisable to consider pregnancy within the first year post transplantation and it is essential that you use some form of contraception. After the first year, if you wish to consider having a baby you should contact the medical staff at the clinic for advice. Many women who have had a successful transplant operation enjoy a normal pregnancy, however you may need to have some of your drugs changed during your pregnancy and also if you want to breastfeed your baby.

**Infertility** - Some patients may experience difficulty in conceiving/ fathering a child. Please bring this to the attention of the medical staff. It is not uncommon and we can refer you for expert help.

**Breast checks** - This should be performed at least monthly and it is best to undertake these tests at a different time from your period. If you are unsure how to perform these tests our medical staff will instruct you. We also have leaflets available. When you examine your breasts you are checking for any swelling, lumps, skin breakdown and discharge from the nipple, remembering to check under your arm as well. Please inform your GP or renal physician if you discover any of the above.

**Smear Tests** - as with all women who are sexually active, it is important to have a smear test every three years.

**Testicular checks** - All men should check at least monthly for any swelling or lumps. If you do not know how to examine yourself, please ask a member of the medical staff who will advise you or we have leaflets available.

**Travel** - There is no reason to prevent you from travelling within the UK when you feel well enough but you should contact one of the medical staff you see regularly in the transplant follow-up clinic for advice. We do not recommend overseas travel in the first six months post transplantation. Before booking any overseas holidays always check that you are fit for travel by asking one of the medical staff in the outpatient clinic. If you are going away you should always take this booklet and sufficient medication with you for your holiday. It is usually a good idea to obtain prescription supplies well in advance so as never to be short of medication. Transplant medications make you more susceptible to skin cancer, so sunscreens and hats should be worn during periods of sun exposure. You will also need to get advice from your GP on vaccinations and antimalarials well before holidays or travel abroad. As you are taking immunosuppressants, it is important that you do not have certain vaccines, called live vaccines, as this is likely to give you the disease the vaccine is designed to protect you against. Other vaccines, called inactivated vaccines are not dangerous to receive, but might not give you full protection.

**Alcohol** - It is possible to drink alcohol within the limits of the National guidelines:

men - 3 - 4 units daily                      women - 2 - 3 units daily

1 glass of wine = 2.3 units                      1 pint of beer - 2.8 units

*Aim to have at least two alcohol-free days a week.*



## Social worker

We appreciate that coming into hospital can be stressful for both patients and their families for a whole variety of reasons. A social worker is available to assist and support you throughout your stay and following your discharge. You may already have met with them during your spell on dialysis.

Information related to Welfare Benefits and a few general guidelines are listed below. If you are in doubt, you should contact your local Benefits Office or speak with one of the social workers.

## What benefits am I entitled to?

**Fares:** If you are in receipt of Income Support, Income Based Job Seeker's Allowance, Income-Related Employment Support Allowance (ESA) or Guarantee Pension Credit, you should be able to claim fares for your trips to hospital. You should ask the ward or clinic staff for a Certificate of Attendance and take this to the hospital cashier, along with proof of benefit, to reclaim your fares.

You may also qualify for help if your income is £15,276 or less and you also get either Child Tax Credit (with or without Working Tax Credit) or Working Tax Credit with the disability element or severe disability element. If you are on a low income but don't get any of these benefits or allowances, you may still be able to claim travel costs through NHS Low Income Scheme. To apply for the NHS Low Income Scheme, you need to fill in form HC1. You can order form HC1 by phoning the NHS Patient Services Helpline on 0845 850 1166 or get it from the RIE Social Work Department. Your form will be assessed and if you're entitled, you'll get a certificate that confirms whether you receive full or partial help with your hospital travel costs.

Unfortunately, fares are not paid for those visiting you. If, however, they receive Income Support, they should approach their local Benefits Agency and request a grant or loan from the Social Fund. If your relatives are finding it financially difficult to visit you, have a word with one of the social workers and it may be possible to get a small grant from one of the hospital or kidney charities.

**Income Support:** If you receive Income Support, you should advise your local Benefits Office of your admission to hospital. Your benefit will continue for six weeks before being reduced.

DLA (Care Component) and Attendance Allowance: These benefits will stop after four weeks in hospital. You should, once again, advise your local Benefits Office of your admission. Carer's Allowance will also be withdrawn at this four-week stage.

## Patient label

I ..... , the patient being assessed for transplantation, have read this booklet and have had the opportunity to discuss the content and other issues important to me with the transplant team.

I am aware that information about me and my transplant will be passed on to relevant healthcare professionals in relation to my care. This information will also be kept on file or a data base in the hospital and with National/International registries. e.g. Organ Donation and Transplantation. This information will be used to demonstrate the results of transplantation as a treatment option and help make the decision about who receives a kidney. The data will be kept in an anonymous form unless it is vital to know the identity of an individual.

Signed: .....

Date: .....

A copy of this completed form will be placed in your transplant medical records.



